



SLEEPLESSNESS (INSOMNIA) AND ITS IMPACT ON THE HUMAN BODY

Ergasheva Munis Muxiddin qizi

munisergasheva45@gmail.com

2nd-year student at the Termez Branch of Tashkent State Medical University

+998970700116

Hasanov Dilmurod Abduvahob o'g'li

dilmurodhasanov1995@gmail.com

Termez Branch of Tashkent State Medical University

Assistant at the Department of Anatomy and Clinical Anatomy

Abstract: This study analyzes the psychological factors contributing to the onset of insomnia and its impact on human physiology. Psychological factors—such as stress, anxiety, depression, emotional strain, and poor sleep habits—are identified as primary causes of sleep disturbances. The effects of insomnia on the body include hormonal imbalances, a weakened immune system, impaired cardiovascular function, and metabolic changes. Furthermore, a decline in cognitive functions—specifically attention, memory, and emotional stability—is noted as a significant consequence of insomnia. The research findings indicate that adhering to sleep hygiene practices and maintaining psychological well-being are crucial for the prevention of insomnia.

Keywords: insomnia, stress, anxiety, depression, psychological factors, sleep disturbance, melatonin, cortisol, physiological changes, immunity, cardiovascular system, metabolism, cognitive functions, emotional instability, sleep hygiene.

Introduction: Sleep is one of the fundamental physiological processes essential for a person's physical and mental health. Stress, anxiety, emotional strain, and an unhealthy lifestyle in modern life are contributing to the widespread prevalence of insomnia. Insomnia not only disrupts one's psychological state but also negatively affects hormonal balance, the immune system, the cardiovascular system, and metabolism. This topic aims to provide a brief overview of the psychological factors behind insomnia and the mechanisms through which it impacts the body's physiology.

Psychological factors in insomnia: Psychological factors play a primary role in the development of insomnia. First and foremost is stress—a state of mental tension arising from the pressures of daily life, demands at work or school, family issues, or financial concerns. Under stress, the nervous system enters a state of heightened activity, preventing the body from engaging in the physiological processes necessary for relaxation. This leads to difficulty falling asleep or frequent awakenings during the night.



Anxiety is one of the most common causes of insomnia. Excessive worry, concern about future events, and overthinking keep the brain in an active state even in the evening. Consequently, sleep quality declines, and transitioning into the deep sleep stage becomes difficult.

Depression is also a significant psychological factor contributing to sleep disturbances. In individuals experiencing depression, the balance of hormones that regulate mood and sleep—such as melatonin and serotonin—is disrupted. This results in irregular sleep patterns, such as waking up too early or lying awake for long periods during the night. Emotional tension—such as an intensification of feelings like anger, resentment, fear, or distress—causes excessive arousal in the nervous system. In such states, it becomes difficult for a person to calm down before falling asleep.

Furthermore, poor sleep habits are closely linked to psychological factors. Staring at the screen of a phone, computer, or television until late at night overloads the brain with information and disrupts its natural preparation for sleep. The blue light emitted by these screens reduces the release of melatonin, thereby throwing off the sleep cycle.

The impact of sleep deprivation on the body's physiology: Sleep deprivation has a profound effect not only on mental health but also on physical well-being. When the body's natural biological rhythms are disrupted, negative physiological changes occur across various systems. Primarily, because sleep serves as the essential period for the body's recovery and regeneration processes, a lack of it leads to numerous functional impairments.

Hormonal imbalance is one of the most significant consequences of sleep deprivation. During the night, the body should produce high levels of the hormone melatonin, which regulates sleep cycles and supports immune function. Sleep deprivation causes a sharp decline in melatonin levels. Furthermore, cortisol—a stress hormone—remains elevated at night instead of peaking in the morning. This disrupts physiological rhythms, leads to overstimulation of the nervous system, and increases the strain on the cardiovascular system.


Another major consequence of sleep deprivation is a weakened immune system. At night, immune cells become active, protecting the body against viruses and bacteria. When sleep is insufficient, this defense system weakens, resulting in an increased susceptibility to illness.

The cardiovascular system also suffers significantly as a result of sleep deprivation. Chronic sleep deprivation raises blood pressure, accelerates the heart rate, and reduces the elasticity of blood vessels. Long-term sleep deficiency can increase the risk of hypertension, ischemic heart disease, and even stroke.

Lack of sleep also negatively affects metabolism. During healthy sleep, the body regulates its energy expenditure. Sleep deprivation, however, decreases insulin sensitivity and disrupts glucose metabolism. This leads to weight gain and an

increased risk of diabetes. In particular, staying awake at night disrupts the balance between appetite-regulating hormones (ghrelin and leptin), which encourages overeating.

A decline in the functioning of the nervous system is also a significant physiological consequence of sleep deprivation. During sleep, neurons in the brain undergo restoration, harmful substances are cleared away, and memories are consolidated. A lack of sleep leads to the accumulation of toxic proteins in the brain, a slowing of reflexes, impaired coordination, and reduced attention.



How Much Sleep Is Enough?

Experts have created guidelines for the amount of sleep needed to maintain optimal mental health, physical health, and emotional well-being. Without this amount of sleep, people begin to accumulate sleep debt and experience the consequences of sleep deprivation. Sleep needs vary across the lifespan but also from person to person. This means that how much sleep an individual needs depends on more than just their age. For example, some people may be naturally long or naturally short sleepers who require more or less than the recommended number of hours to wake up feeling rested.

How much sleep a person needs also depends on their health and typical daily activities. In the short-term, the need for sleep is temporarily increased after demanding activities, when a person is sick, or when recovering from a period of sleep deprivation.

Additionally, avoiding sleep deprivation is about more than just spending enough hours in bed. Healthy and restorative rest also depends on the quality of sleep.



So even if a person gets the right amount of hours of sleep, they may still be sleep deprived if their sleep quality is reduced from waking up too often at night.

Age Group	Age Range	Recommended hours of sleep per 24 hours
Infant	4 to 12 months	12 to 16 hours (including naps)
Toddler	1 to 2 years	11 to 14 hours (including naps)
Preschool	3 to 5 years	10 to 13 hours (including naps)
School-age	6 to 12 years	9 to 12 hours
Teen	13 to 18 years	8 to 10 hours
Adult	18 years and older	7 to 9 hours

Preventing Sleep Issues

To reduce the risk of sleep deprivation, it is important to take steps to improve sleep hygiene.

Make sleep a priority: Prioritize sleep health by creating a comfortable sleep environment and keeping a consistent sleep schedule. This means going to sleep and waking up at around the same time each day and avoiding the temptation to stay up later or sleep in on the weekends.

Combat stress: To combat bedtime stress, give yourself plenty of time to wind down from the day. Use this time to listen to calming music, stretch, or write in a journal. Boost your ability to relieve stress by trying out new relaxation techniques and seeing what helps the most.

Time your light exposure: Ambient light can signal to the body whether it is time to be awake or prepare for sleep, so be intentional about light exposure. Try to get at least 30 minutes of sunlight exposure during the day and then dim or turn off lights in the evening. Shut off phones, TVs, and computers at least an hour before bed.

Watch your caffeine intake: Caffeine can linger in the body for eight or more hours, so consuming caffeine in the afternoon may affect how long it takes to doze off at bedtime.

Nap wisely: Although they cannot replace quality nighttime rest, naps can be a helpful tool to improve daytime alertness. If naps are too long or poorly timed, though, they can make it more difficult to fall asleep in the evening. Adults should aim for naps that are no longer than 20 minutes and should avoid napping in the late afternoon.

Stay active: Regular exercise can make it easier to get to sleep at bedtime. Try to get at least 30 minutes of physical activity every day, but it is best to avoid highly strenuous exercise too close to bedtime.



In conclusion, sleep deprivation is a complex condition, prevalent in modern society and closely linked to both psychological and physiological factors. Stress, anxiety, depression, emotional strain, and an unhealthy lifestyle disrupt the sleep process and constitute the primary psychological causes of sleep deprivation. Such conditions overstimulate the central nervous system, hindering the body's mechanisms for relaxation and recovery. The physiological impact of sleep deprivation on the body is extensive, leading to serious consequences such as hormonal imbalances, a weakened immune system, impaired cardiovascular function, metabolic changes, and a decline in nervous system activity. Furthermore, prolonged sleep deprivation increases the risk of chronic diseases and significantly diminishes overall quality of life and productivity.

References

1. Ramar K, Malhotra RK, Carden KA, et al. Sleep is essential to health: an American Academy of Sleep Medicine position statement. *J Clin Sleep Med*. 2021;17(10):2115-2119. doi: 10.5664/jcsm.9476
2. Buysse DJ. Sleep health: can we define it? Does it matter? *Sleep*. 2014;37(1):9-17. doi: 10.5665/sleep.3298
3. Lim DC, Najafi A, Afifi L, et al. The need to promote sleep health in public health agendas across the globe. *Lancet Public Health*. 2023;8(10):e820-e826. doi: 10.1016/S2468-2667(23)00182-2
4. Goldstein AN, Greer SM, Saletin JM, Harvey AG, Nitschke JB, Walker MP. Tired and apprehensive: anxiety amplifies the impact of sleep loss on aversive brain anticipation. *J Neurosci*. 2013;33(26):10607-10615. doi: 10.1523/JNEUROSCI.5578-12.2013
5. Watson NF, Badr MS, Belenky G, et al. Recommended amount of sleep for a healthy adult: a joint consensus statement of the American academy of sleep medicine and sleep research society. *Sleep*. 2015;38(6):843-844. doi: 10.5665/sleep.4716
6. Hirshkowitz M, Whiton K, Albert SM, et al. National sleep foundation's sleep time duration recommendations: methodology and results summary. *Sleep Health*. 2015;1(1):40-43. doi: 10.1016/j.sleh.2014.12.010
7. Mukherjee S, Patel SR, Kales SN, et al. An official American thoracic society statement: the importance of healthy sleep. Recommendations and future priorities. *Am J Respir Crit Care Med*. 2015;191(12):1450-1458. doi: 10.1164/rccm.201504-0767ST
8. Gallicchio L, Kalesan B. Sleep duration and mortality: a systematic review and meta-analysis. *J Sleep Res*. 2009;18(2):148-158. doi: 10.1111/j.1365-2869.2008.00732.x