



EVALUATION OF NEWBORN INFANTS ACCORDING TO THE APGAR SCORING SYSTEM

Zebo Ravshanovna Xudoyqulova

zeboxudoyqulova1980@gmail.com

Termez University of Economics and Service, Faculty of Medicine

Lecturer, Department of Medical Clinical Sciences

Jasur Baxtiyarovich Saidov

<https://orcid.org/0009-0007-1601-1838>

jasur_saidov@tues.uz

Termez University of Economics and Service, Faculty of Medicine

Lecturer, Department of Medical Clinical Sciences

Ismatullayeva Muxlisa

Student, Faculty of Medicine

Termez University of Economics and Service, Termez, Uzbekistan

Annotation: The APGAR scoring system is a simple and effective method used worldwide to assess the physical condition of newborn infants immediately after birth. Developed by Dr. Virginia Apgar in 1952, this scoring system evaluates five essential clinical parameters: heart rate, respiratory effort, muscle tone, reflex irritability, and skin coloration. The APGAR score provides rapid information about the newborn's adaptation to extrauterine life and helps healthcare professionals identify infants requiring immediate medical intervention. This article examines the significance, methodology, and clinical applications of APGAR assessment in neonatal care. The role of APGAR scoring in improving neonatal outcomes and reducing the risk of complications is also discussed.

Keywords: APGAR score, newborn infant, neonatal assessment, neonatal care, birth evaluation, respiratory effort, heart rate, reflex response, muscle tone, neonatal health.

Introduction



The transition from intrauterine to extrauterine life is one of the most critical periods in human development. Immediately after birth, newborn infants must rapidly adapt to significant physiological changes, including independent breathing, circulatory adjustments, and temperature regulation. Early assessment of a newborn's condition is therefore essential for ensuring timely medical care and reducing neonatal morbidity and mortality. The APGAR scoring system, introduced by anesthesiologist Virginia Apgar, has become one of the most widely used tools for evaluating neonatal well-being. The assessment is typically performed at one and five minutes after birth and may be repeated when necessary. The score is based on five observable clinical signs: heart rate, respiratory effort, muscle tone, reflex irritability, and skin color. Each parameter is assigned a score ranging from 0 to 2, resulting in a total score between 0 and 10. APGAR scoring provides a rapid and standardized method for determining the newborn's physiological status and need for immediate intervention. Although it is not designed to predict long-term neurological outcomes, it remains a valuable indicator of the infant's condition during the first minutes of life. The widespread use of the APGAR system has contributed significantly to advances in neonatal care and the improvement of newborn survival rates worldwide. This article aims to review the principles of APGAR assessment, its clinical significance, and its role in the early identification and management of neonatal complications.

Materials and Methods

This study was conducted to evaluate the effectiveness and clinical importance of the APGAR scoring system in assessing the health status of newborn infants immediately after birth. The research was based on a review of scientific literature, neonatal care guidelines, and clinical observations obtained from maternity and neonatal units. The APGAR score was assessed at one and five minutes after birth using five standard criteria: heart rate, respiratory effort, muscle tone, reflex irritability, and skin coloration. Each criterion was assigned a score of 0, 1, or 2 points, resulting in a total score ranging from 0 to 10. Newborns were classified into three groups according to their scores: severe distress (0–3 points), moderate difficulty (4–6 points), and good condition (7–10 points). Data analysis focused on the relationship between APGAR scores and the immediate health outcomes of newborns. Particular attention was given to the need for resuscitation, respiratory support, and admission to neonatal intensive care units.

Results

The findings indicated that the APGAR scoring system provides a rapid and reliable assessment of neonatal adaptation following birth. The majority of newborn infants achieved APGAR scores between 7 and 10, indicating satisfactory physiological adjustment to extrauterine life. Infants with



APGAR scores of 4–6 frequently required additional observation and supportive care, including oxygen therapy and monitoring of vital signs. Newborns with scores between 0 and 3 demonstrated significant physiological compromise and often required immediate resuscitation measures and intensive medical management. A substantial improvement in APGAR scores was observed between the first and fifth minutes after birth in many infants, reflecting successful adaptation and effective neonatal interventions. The five-minute APGAR score showed a stronger association with short-term neonatal outcomes than the one-minute score. The study also revealed that premature infants and newborns delivered after complicated pregnancies were more likely to have lower APGAR scores compared to healthy full-term infants.

Discussion

The APGAR scoring system remains one of the most valuable tools in neonatal medicine due to its simplicity, speed, and clinical usefulness. The results of this study confirm that APGAR assessment provides essential information regarding the newborn's immediate physiological condition and the effectiveness of adaptation to life outside the womb. The ability to identify infants experiencing respiratory distress, circulatory instability, or neurological depression within the first minutes after birth allows healthcare professionals to initiate prompt interventions. Early recognition and treatment significantly reduce the risk of neonatal complications and improve survival rates. Although APGAR scoring is highly effective for immediate assessment, it should not be used as the sole predictor of long-term neurological or developmental outcomes. Additional clinical examinations and diagnostic investigations are often required to obtain a comprehensive evaluation of neonatal health. The findings support the continued use of the APGAR system as a standard component of neonatal care worldwide. Its universal application facilitates communication among healthcare providers and contributes to the consistent assessment and management of newborn infants. Overall, APGAR scoring remains an indispensable method for evaluating newborn health, guiding clinical decision-making, and ensuring the timely delivery of life-saving interventions during the critical postnatal period.

Conclusion

The APGAR scoring system is a simple, rapid, and effective method for evaluating the condition of newborn infants immediately after birth. By assessing five essential physiological parameters—heart rate, respiratory effort, muscle tone, reflex irritability, and skin coloration—the APGAR score provides valuable information about a newborn's adaptation to extrauterine life. The findings of this study demonstrate that APGAR assessment plays a crucial role in the early identification of infants requiring medical attention and resuscitative measures. Newborns with low APGAR scores are at



increased risk of respiratory and circulatory complications, whereas higher scores generally indicate successful physiological adaptation. Although the APGAR score is not intended to predict long-term developmental outcomes, it remains an indispensable tool in neonatal medicine. Its ease of use, reliability, and universal acceptance make it an essential component of newborn assessment worldwide. Continuous application of the APGAR scoring system contributes to improved neonatal care, timely clinical interventions, and better health outcomes for newborn infants.

References:

1. Apgar V. A proposal for a new method of evaluation of the newborn infant. *Current Researches in Anesthesia and Analgesia*. 1953;32(4):260–267.
2. American Academy of Pediatrics, American College of Obstetricians and Gynecologists. The Apgar Score. Committee Opinion No. 644. *Pediatrics*. 2015;136(4):819–822.
3. Cunningham F.G., Leveno K.J., Bloom S.L., et al. *Williams Obstetrics*. 26th ed. New York: McGraw-Hill Education; 2022.
4. Kliegman R.M., St. Geme J.W. *Nelson Textbook of Pediatrics*. 22nd ed. Philadelphia: Elsevier; 2023.
5. Martin R.J., Fanaroff A.A., Walsh M.C. *Fanaroff and Martin's Neonatal-Perinatal Medicine*. 12th ed. Elsevier; 2025.
6. World Health Organization. *Recommendations on Newborn Health: Guidelines Approved by WHO*. Geneva: WHO; 2017.
7. Finster M., Wood M. The Apgar score has survived the test of time. *Anesthesiology*. 2005;102(4):855–857.
8. Casey B.M., McIntire D.D., Leveno K.J. The continuing value of the Apgar score for assessing newborn infants. *New England Journal of Medicine*. 2001;344(7):467–471.
9. Papile L.A. The Apgar score in the 21st century. *New England Journal of Medicine*. 2001;344(7):519–520.
10. World Health Organization. *Standards for Improving Quality of Maternal and Newborn Care in Health Facilities*. Geneva: WHO; 2016.